


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/yexcr

Review Article

Regulation of cell adhesion and migration by cell-derived matrices



Matthew L. Kutys*, Andrew D. Doyle, Kenneth M. Yamada

Laboratory of Cell and Developmental Biology, National Institute of Dental and Craniofacial Research, United States

ARTICLE INFORMATION

Article Chronology:

Received 18 March 2013

Received in revised form

30 May 2013

Accepted 31 May 2013

Available online 7 June 2013

Keywords:

Cell-derived matrix (CDM)

Extracellular matrix

3D cell migration

Matrix adhesions

Fibrillar topography

Matrix elasticity

ABSTRACT

Three-dimensional *in vitro* extracellular matrix models provide a physiological alternative to regular two-dimensional cell culture, though they lack the full diversity of molecular composition and physical properties of whole-animal systems. Cell-derived matrices are extracellular matrices that are the product of matrix secretion and assembly by cells cultured at high density *in vitro*. After the removal of the cells that produced the matrix, an assembled matrix scaffold is left that closely mimics native stromal fiber organization and molecular content. Cell-derived matrices have been shown to impart *in vivo*-like responses to cells cultured in these matrices. In this review, we focus on mechanisms through which the distinct molecular and topographical composition of cell-derived matrices directs cellular behavior, specifically through regulation of cell-matrix adhesions and subsequent contributions to the process of cell migration.

Published by Elsevier Inc.

Contents

Introduction	2435
Biochemical and biophysical characterization of CDMs	2435
Cell-derived matrices regulate adhesion composition and stability	2436
Fibrillar topography dictates cell migration in cell-derived matrix.	2437
Applications of cell-derived matrix.	2438
Conclusion	2438
Acknowledgments	2438
References	2438

*Correspondence to: National Institutes of Health, National Institute of Dental and Craniofacial Research, Building 30, Room 426, 30 Convent Dr MSC 4370, Bethesda, MD 20892-4370, United States. Fax: +1 301 402 0897.

E-mail addresses: kutysml@mail.nih.gov (M.L. Kutys), kyamada@mail.nih.gov (K.M. Yamada).

Introduction

The use of 3D *in vitro* models to bridge the gap between cell culture and *in vivo* systems permits greater understanding of the physical and chemical interactions between cells and the 3D extracellular matrix (ECM). Cell-derived matrices (CDMs) provide a more complex, physiological alternative to purified 3D matrix protein scaffolds, such as polymerized type I collagen or fibrin gels; CDMs consist of a heterogeneous mixture of proteins, proteoglycans and growth factors similar to the native stromal environment [1]. CDMs are naturally produced by the continued secretion and organization of ECM components into a 3D matrix when certain cell types are cultured at high density *in vitro*. Within several (3–10) days, a thick 3D matrix is generated that can be denuded of cells to yield a 3D ECM substrate [2]. Fibroblasts are responsible for *in vivo* stromal ECM deposition and maintenance and are commonly used for CDM generation, though 3D CDMs have been produced from endothelial, epithelial, stem cell, and cancer cell cultures [3–6].

Biochemical and biophysical characterization of CDMs

Fibroblast-generated CDM is a heterogeneous fibrous matrix, consisting primarily of a meshwork of linear fibronectin fibrils that can be oriented in parallel or more random in organization (Fig. 1, magenta fibronectin fibers). Additional matrix proteins such as collagen I and IV, perlecan, tenascin-C, hyaluronic acid, and heparan sulfate proteoglycans are present in lower abundance, as well as sequestered growth factors [3,7]. This diversity and spatial heterogeneity of CDM components mimic what is found in *in vivo* matrix, providing physiological properties not commonly found in traditional polymerized or synthetic 3D *in vitro* scaffolds. Because CDM is generated by the secretion and assembly of matrix fibers from layers of confluent cells *in vitro*, its topography consists of arrays of fibronectin fibers that are stacked to an approximate depth of 5–20 μm [5,8–10]. These fibers have an average inter-fiber spacing of approximately 1–2 μm [9], resulting in larger internal spaces in comparison to other 3D ECM models. This distinct matrix architecture provides an advantage for microscopy by minimizing light scatter through the matrix and additionally ensures relatively constant nutrient delivery and physical characteristics throughout the CDM model.

One concern raised about most 3D *in vitro* matrix models, whether comprised of a purified protein or CDM, is the physiological relevance of its matrix stiffness, particularly regions in close proximity to a rigid underlying glass or plastic surface (>1 GPa). This issue of local matrix properties is relevant biologically, since ECM stiffness is sensed by cells through cell-matrix adhesions and can alter intracellular signaling pathways, such as by Rho GTPases, leading to changes in migration, differentiation and proliferation [11–13]. Atomic force microscopy measurements of CDM have reported Young's modulus in the range of 200–600 Pa, which is comparable to reconstituted basement membrane (175 Pa), 3D collagen gels (concentration dependent, 15–1000 Pa), and dermal tissue explants (>300 Pa) [8,9,13,14]. Despite their relatively shallow depth and proximity to an artificial, rigid surface, CDMs are a physiologically-compliant




Fig. 1 – Fibroblast migration in 3D cell-derived matrix. Left: maximum projection confocal image of a human foreskin fibroblast (HFF) embedded within a matrix containing oriented CDM fibronectin fibrils (magenta, anti-fibronectin immunostaining). Cytoskeletal architecture visualized by staining for actin (blue, phalloidin) and paxillin-containing focal adhesions (yellow, anti-paxillin). Right: maximum projection image of focal adhesion profile of paxillin-containing adhesions during migration in CDM. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

matrix, allowing for flexibility and malleability similar to *in vivo* ECM. An additional physical parameter that has recently been identified as a regulator of cell migration and signaling is the elastic behavior of the matrix, *i.e.*, non-linear or linear elasticity. In contrast to non-linearly elastic 3D collagen matrices, CDMs are linearly elastic and do not undergo strain stiffening under increasing force, perhaps due to its distinct composition and crosslinking between fibers compared to meshwork collagen matrices. Using CDM, Petrie et al. demonstrated that fibroblasts have the ability to discern the elastic behavior of the ECM and accordingly switch modes of migration [14].

Besides possessing biochemical and physical characteristics that mimic *in vivo* matrices, CDM is by nature an *in vitro* system and thus more accessible to experimental manipulation. The composition of the medium used for culturing cells during CDM production can be adjusted to alter matrix composition, thickness, and topography. For example, the addition of ascorbic acid to culture media will increase the collagen content of the CDM and addition of epidermal growth factor (EGF) and basic fibroblast growth factor (bFGF) will increase overall ECM production and matrix thickness [8,9]. Genetic ablation and RNA interference analyses of the cultured fibroblasts during CDM production can permit direct

examination of the pathways involved in ECM synthesis, deposition, and organization. Recently, Goetz et al. discovered a role for caveolin-1 in the topographical assembly and remodeling of 3D microenvironments by examining the organization and compliance of CDM fibronectin fibers derived from wild-type and caveolin-1 knockout fibroblasts [15]. Additionally, post-synthesis chemical manipulation of CDM by crosslinking the matrix or by treating with trypsin can modulate the stiffness and elastic behavior of the CDM [14]. Taken together, CDM provides an *in vitro* 3D model that closely mimics the chemical and physical heterogeneity found *in vivo*, but which also allows for experimental manipulation in composition and physical properties to test hypotheses in ways that are not possible *in vivo*.

Cell-derived matrices regulate adhesion composition and stability

It is well known that mesenchymally derived fibroblasts require attachment or adhesion to the surrounding extracellular matrix for cell migration as well as maintenance of cell homeostasis and survival. Other cell types, such as many cancer cell lines and hematopoietic cells can not only migrate, but can survive in the absence of cell-matrix attachment [16]. In this section, we will focus solely on cell-CDM adhesions made by fibroblasts and other anchorage-dependent cell types.

Cell-matrix interactions at cell adhesions not only provide physical support for cell migration, but can also act as outside-in signaling hubs that initiate or modulate a wide range of intracellular signaling cascades. Linkage to the surrounding matrix is often initiated by integrins, a group of heterodimeric transmembrane proteins consisting of 13 α and seven β subunits, whose different combinations dictate their ECM specificity [17]. While multiple pairs of α and β integrin subunits can mediate adhesion to the same type of ECM molecule, the strength of their association can vary greatly. For example, both $\alpha v \beta 3$ and $\alpha 5 \beta 1$ can bind to fibronectin, the primary component of 3D CDMs, yet the adhesion strength of $\alpha 5 \beta 1$ -containing adhesions can be six-fold higher [18]. Within 3D CDMs, the high quantity of fibronectin tends to favor $\alpha 5 \beta 1$ associated attachments, yet Chinese hamster ovary (CHO) cells that lack the $\alpha 5$ subunit can still attach and become imbedded in the matrix through $\alpha v \beta 3$ -matrix interactions [19]. A key connection between integrins and the surrounding ECM is through the β subunit, though the α subunits are also essential for function. A number of integrins can bind to short peptide sequences in matrix proteins, such as the arginine-glycine-aspartic acid (RGD) sequence found on fibronectin, vitronectin, and other ECM proteins [20].

The cytoplasmic tails of integrins interact with numerous intracellular proteins in large multimolecular complexes at cell adhesions. These proteins vary widely in function, from mechanosensitive proteins such as vinculin, to signaling molecules like Src, to cytoskeletal adapter proteins such as paxillin (Fig. 1, yellow adhesions). These proteins initially localize or accumulate at clustered integrins to form small foci that can become directly linked to the actin cytoskeleton. It is at these cellular adhesions that actomyosin contractility can be propagated to the external environment. Hence, cellular adhesions are the sites of mechanotransduction between the cell and its surroundings. We use the term cell adhesions as a loose general classification that include

nascent adhesions, focal complexes, focal adhesions, fibrillar adhesions, and 3D matrix adhesions; each can be considered a distinct subset of adhesions with different molecular components and associations with the actin cytoskeleton [21].

The majority of adhesion studies have been performed on 2D rigid glass surfaces, often coated with an adsorbed globular ECM molecule, such as fibronectin. In general, 2D cell-ECM adhesions proceed through a maturation process; that is they grow in size, increase the number of proteins that associate within them, and progress in order from focal contacts through fibrillar adhesions [22]. While both nascent adhesions and focal complexes have been observed and well-characterized on 2D substrates, they have not been identified within 3D CDM to date. This is likely due to their small size, relatively short longevity (under 90 s), and absence of requirement for actin association [23]. However, the absence of nascent adhesions in 3D CDMs suggests that nearly all develop into actin-linked, contractility-dependent adhesions [24]. 3D matrix adhesions (3D-MA) formed by cells with 3D matrix fibrils have characteristics of both focal (FA) and fibrillar (FX) adhesions [10]. More than a dozen years ago, Cukierman et al. [10] determined several key protein components of 3D-MA in CDM including: activated $\beta 1$ integrin, $\alpha 5$ integrin, α -actinin, actin, talin, tensin, vinculin, focal adhesion kinase (FAK) and paxillin. Interestingly, FAK in 3D-MA is not phosphorylated at tyrosine 397, similar to fibrillar adhesions in 2D, while 2D FAs have highly phosphorylated FAK. Phosphorylation of FAK at this site has been correlated with rigidity sensing on 2D rigid surfaces, and is down-regulated on soft 2D substrates or in 3D collagen gels [25,26]. Additionally, even though $\beta 3$ integrin was plentiful in 2D FAs, it is absent from 3D-MA structures in CDM [21].

3D CDMs have been utilized to elucidate mechanisms of cancer cell invasion. Deakin and Turner have shown that the adapter proteins hic-5 and paxillin are crucial for cancer cell attachment or for mediating 3D-MA adhesion dynamics in 3D CDMs, respectively [27]. Recently, proteomic analysis has identified greater than 700 different proteins and/or macromolecular complexes that reside within myosin II-dependent FAs formed on 2D rigid surfaces with 283 known to be a part of the integrin adhesome [28]. While this feat has yet to be accomplished for cells embedded within 3D CDMs, it seems likely that many FA-associated proteins will be found in 3D-MAs.

The coupling that occurs within cellular adhesions links integrins and the actin cytoskeleton. This integration site has been dubbed the molecular clutch, where relatively immobile integrins are indirectly coupled to rapidly flowing actin through binding interactions involving the numerous cell adhesion-associated proteins discussed above. It is now known that cellular adhesions are highly dynamic structures that assemble, grow or mature, become stable, and finally disassemble [24] with the latter three stages being highly dependent on actin-based contractility. Hence, nascent adhesions that are contractility-independent have a short lifespan or stability compared to contractility dependent FAs. Recent evidence suggests that the coupling between 3D-MAs to CDM is relatively high, showing a three-fold increase in length of time that these adhesions exist compared to 2D FAs [23]. One possible explanation is the high concentration of fibronectin within 3D CDMs: increasing ligand density on 2D surfaces leads to a decrease in FA turnover [29]. In addition, the fibrillar structure of fibronectin may also be a regulating factor. When fibronectin is physically stretched during the process of

fibrillogenesis, interactions occur between multiple cryptic cysteine residues within several fibronectin-type III (FN III) repeats between molecules, which has been shown to increase the ability of fibronectin to withstand forces compared to single molecules alone [30]. These ideas suggest that adhesion stability within 3D CDMs may be intrinsically related to the concentration of ECM proteins (specifically fibronectin) and their fibrillar nature.

One very important feature of 3D-MAs that is not normally found in 2D FAs is the abundance of $\alpha 5\beta 1$ integrin localizing to adhesion sites. This localization is likely due to the high concentrations fibronectin in 3D CDMs. Hence, research investigating the interactions between integrins and fibronectin, regardless of ECM dimensionality, can contribute to our understanding of cellular adhesion to 3D CDMs. For example, increasing the concentration of fibronectin (from ~ 10 to $50 \mu\text{g/ml}$) on 2D surfaces results in the appearance of $\alpha 5$ integrin in 2D FAs (A.D.D., unpublished results), which could explain the high amounts of this integrin in 3D-MAs interacting with thick fibrils of fibronectin. In contrast, $\alpha V\beta 3$ is normally present in 2D FAs at low fibronectin concentrations of $1\text{--}10 \mu\text{g/ml}$. Chemical crosslinking of 2D FN-coated dishes also produces a similar migration of $\alpha 5$ into FAs [31]. The amino terminal domain of the $\alpha 5$ subunit is known to bind directly to fibronectin at the FN III-9 repeat, also known as the synergy site [32]. Although the RGD-binding site on $\beta 1$ is the most crucial for binding ligand, the synergy site considerably promotes binding to fibronectin [33], especially when a cell is under tension. Friedland et al. [34] demonstrated that when bound to fibronectin, elevating cytoskeletal or ECM tension results in a switch of the $\alpha 5\beta 1$ integrin from a relaxed to a tension state. This switch is associated with engagement of $\alpha 5$ with the synergy site. Mutating the synergy site resulted in weakening of the integrin–fibronectin linkage, suggesting that it is required for increasing ligand–receptor bond strength and likely the stabilization of cellular adhesions.

In summary, the adhesion components that constitute 3D-MAs show similarities with 2D FAs and FXs [23], but how these components are utilized for intracellular signaling remains to be determined. The major differences in molecular composition between these subtypes of cell adhesions are likely due to the ECM itself, where high concentrations of organized fibrillar fibronectin promote $\alpha 5\beta 1$ integrin association, increasing cell adhesiveness and likely altering outside-in signaling events.

Fibrillar topography dictates cell migration in cell-derived matrix

Cell migration studies in 3D have achieved prominence due to the stark differences in morphology, motility, adhesion, and signaling observed in the same cell type by simply altering matrix dimensionality. These distinct phenotypes result from differences between 2D and 3D migratory environments, including presence of dorsal and ventral matrix adhesions, fibrillar topography of the matrix proteins, and steric inhibition of spreading and migration by matrix fibers and porosity [1,35]. As discussed below, cell migration in CDM is regulated not only by chemical information (molecular composition), but particularly the topographical cues provided by the physical organization of the ECM ligands to which the cells adhere.

In oriented 3D CDMs, such as those generated by primary human fibroblasts that grow in strongly parallel patterns, cells adhere to the linear arrays of fibronectin fibrils, rapidly align and polarize along the fiber, and migrate in a high directional manner. This elongated, uniaxial migratory phenotype mimics the behavior of fibroblasts *in vivo* migrating along stromal fibrils [10,36]. Unlike migration in 3D fibrous matrices such as collagen or fibrin, fibroblasts do not significantly remodel these highly organized CDM fibers. This finding is potentially explained by the inherent linear topography of the fibronectin fibrils generated during CDM production by parallel arrays of cells and the large interfiber spacing ($\geq 2 \mu\text{m}$) [9] of CDM in comparison to the dense lattices of fibrous collagen and fibrin gels. Interestingly, this elongated, spindle-like morphology of cells in 3D CDM contrasts strikingly with the shapes of fibroblasts migrating on 2D fibronectin or on the 2D upper surface of the CDM, both of which are characterized by broad lamellipodia, lower migration rates, and decreased directional persistence in comparison to 3D CDM [10,36]. Recent studies have tested the hypothesis that the fibrillar topography of 3D CDM is the driving mechanism behind this phenotypic switch in migration by mimicking 3D matrix fibrils with 1D fibronectin-coated micropatterned lines. By generating $1.5 \mu\text{m}$ wide fibronectin-coated lines, either as single lines or multiple two micron-spaced arrays, the ligand-independent, rapid uniaxial migratory phenotype observed in 3D CDM was recapitulated on the 1D lines. Examining fibroblasts in 1D and 3D CDM revealed that in comparison to 2D migration, the processes of protrusion, cell translocation, and tail retraction are highly efficiently coordinated in 1D and 3D. Additionally, oriented 1D and 3D matrices provided cues to promote linear actomyosin and microtubule networks, leading to the uniaxial phenotype as well decreased levels of Rac1, which had been previously demonstrated to control directional persistence [36,37]. Taken together, these studies showed 3D CDM migration is highly dependent on physical cues from the linear, fibrillar topography to regulate cell behavior.

Another significant observation made in fibroblasts migrating in 3D CDM is the dependence on myosin II contractility for efficient migration in comparison to 2D. Inhibition of myosin II ATPase activity by the drug blebbistatin produced a 60% decrease in cell velocity during migration in 3D CDM, but in contrast led to a slight increase in velocity and protrusive activity in 2D, indicating a requirement for contractile forces in 3D CDM [23,37]. Leading edge analysis showed a dependence on the spatial distribution of cell adhesions: during fibrillar migration, adhesions formed within one micron of the leading edge and their assembly rates matched the rate of leading edge protrusion. Mechanistically, cell adhesions in 3D CDM are stabilized for prolonged periods as indicated by low turnover rates, and actin retrograde flow is significantly decreased, further demonstrating enhanced coupling between the molecular clutch guiding protrusion and adhesion. Genetic ablation of myosin IIA or IIB isoforms revealed that myosin IIA is critical for maintaining the contractile requirements for efficient migration in CDM. Loss of myosin IIA results in both a loss of the force that is applied to the fibrillar environment and loss of adhesion stability observed during CDM migration [23,37]. Therefore, the physically restricting cues of 3D CDM fibrils require cellular contraction through myosin IIA to allow for increased mechanical coupling and efficient, rapid migration.

Recently, a novel mode of fibroblast motility was identified in both *in vivo*-like dermal explant and CDM models. Originally observed as a distinct population of fibroblasts migrating in CDM, this lobopodial form of migration is defined by the absence of the characteristic lamellipodia at the leading edge of the cell observed during mesenchymal cell migration. Instead, the leading edge of the cell is characterized by a blunt, cylindrical protrusion that is devoid of lamellipodial actin and cortactin; in contrast, blebs appear at the sides of the cell. Interestingly, unlike canonical lamellipodia-based migration, Rac1 and Cdc42 activity are no longer polarized toward the leading edge of the cell. Instead, lobopodial migration is driven by RhoA signaling to downstream effectors Rho-associated protein kinases (ROCK) and myosin II. Through experimental modulation of CDM, lobopodia-based migration was found to be dictated by the linear elasticity of CDM, while lamellipodial-based migration was associated with nonlinear elastic matrix environments, such as polymerized 3D collagen or trypsin-treated CDM [14]. These findings demonstrate that similar to *in vivo* situations, 3D CDM is able to trigger migratory plasticity within the same cell type. Further insight into the molecular mechanisms guiding these migratory switches will be crucial for understanding the processes driving cellular migration through heterogeneous, native stromal tissue.

Applications of cell-derived matrix

CDMs have been shown to impart *in vivo*-like responses to cells cultured inside the matrix. Recent studies have also shown that the CDM produced by cells effectively recapitulates the distinct *in vivo* stromal characteristics of the parental environment of the cells generating that matrix. *In vivo* comparisons of normal and tumor-associated stroma have shown different stromal fiber organization and content, fibroblast morphology, and gene expression. CDM generated from fibroblasts derived from tumor-associated stroma matched the architecture and content observed *in vivo* [6,7,10]. Additionally, tumor-associated CDM was necessary and sufficient to promote desmoplastic differentiation of normal fibroblasts as well as to trigger invasive behavior of breast cancer cells [7,38], indicating that these matrices contain the necessary physical and molecular cues to induce transformation events observed *in vivo* during stromatogenesis. These matrix-encoded signals are also found using *in vitro* models for understanding ECM roles during developmental processes. Studies using CDM generated from young mesenchymal stem cells proved they were able to restore the replication and osteogenic functions of aged mesenchymal stem cells [4]. Fibroblast CDM has also been shown to promote morphogenesis and capillary-like tubule formation of endothelial cells *in vitro* [3,39]. These results with different types of CDMs provide intriguing new approaches to study distinct *in vivo* ECM effects on a variety of cellular processes in a controlled *in vitro* setting.

Conclusion

Many discoveries from *in vivo* and 3D *in vitro* analyses of cellular processes have generated a need for specific, physiologically relevant *in vitro* ECM models. Different types of CDMs can provide the complex composition, stiffness, and topographical

architecture found in native stromal ECM. Cells cultured in CDMs adopt the morphological and migratory phenotypes commonly observed *in vivo*. Compared to traditional 2D migration, cells in 3D CDM can utilize distinct adhesive components and engage the contractile machinery more efficiently, driving persistent, rapid migration. This migratory behavior is directly related to the restrictive fibrillar topography of CDM. These approaches provide insight into the mechanisms driving fibroblast and other mesenchymal cell migration in native stroma. With increasing evidence that CDMs can retain the molecular cues of native stromal tissue, CDMs provide robust *in vitro* models capable of analyzing complex physiological processes in an *ex vivo* setting. The same transfection, RNAi, genetic, and immunological tools used so effectively for understanding cellular functions in regular cell culture can be applied directly to cells growing in CDMs. These approaches should permit more rapid dissection of molecular mechanisms that are difficult to study *in vivo*, and provide new insights into the regulation of cellular functions by the ECM.

Acknowledgments

We thank Ryan Petrie and Will Daley for critical reading of the manuscript. Supported by the Intramural Research Program of the NIDCR, NIH.

REFERENCES

- [1] K.M. Yamada, E. Cukierman, et al., Modeling tissue morphogenesis and cancer in 3D, *Cell* 130 (2007) 601–610.
- [2] K.M. Hakkinen, J.S. Harunaga, A.D. Doyle, K.M. Yamada, et al., Direct comparisons of the morphology, migration, cell adhesions, and actin cytoskeleton of fibroblasts in four different three-dimensional extracellular matrices, *Tissue Eng. Part A* 17 (2011) 713–724.
- [3] P.A. Soucy, L.H. Romer, et al., Endothelial cell adhesion, signaling, and morphogenesis in fibroblast-derived matrix, *Matrix Biol.* 28 (2009) 273–283.
- [4] Y. Sun, W. Li, Z. Lu, R. Chen, J. Ling, Q. Ran, R.L. Jilka, X.D. Chen, et al., Rescuing replication and osteogenesis of aged mesenchymal stem cells by exposure to a young extracellular matrix, *Faseb J.* 25 (2011) 1474–1485.
- [5] C.M. Zwolinski, K.S. Ellison, N. Depaola, D.M. Thompson, et al., Generation of cell-derived three dimensional extracellular matrix substrates from two dimensional endothelial cell cultures, *Tissue Eng. Part C Methods* 17 (2011) 589–595.
- [6] R.M. Quiros, M. Valianou, Y. Kwon, K.M. Brown, A.K. Godwin, E. Cukierman, et al., Ovarian normal and tumor-associated fibroblasts retain *in vivo* stromal characteristics in a 3-D matrix-dependent manner, *Gynecol. Oncol.* 110 (2008) 99–109.
- [7] M.D. Amatangelo, D.E. Bassi, A.J. Klein-Szanto, E. Cukierman, et al., Stroma-derived three-dimensional matrices are necessary and sufficient to promote desmoplastic differentiation of normal fibroblasts, *Am. J. Pathol.* 167 (2005) 475–488.
- [8] J.E. Ahlfors, K.L. Billiar, et al., Biomechanical and biochemical characteristics of a human fibroblast-produced and remodeled matrix, *Biomaterials* 28 (2007) 2183–2191.
- [9] P.A. Soucy, J. Werbin, W. Heinz, J.H. Hoh, L.H. Romer, et al., Microelastic properties of lung cell-derived extracellular matrix, *Acta Biomater.* 7 (2011) 96–105.
- [10] E. Cukierman, R. Pankov, D.R. Stevens, K.M. Yamada, et al., Taking cell-matrix adhesions to the third dimension, *Science* 294 (2001) 1708–1712.

- [11] J.N. Heck, S.M. Ponik, M.G. Garcia-Mendoza, C.A. Pehlke, D.R. Inman, K.W. Eliceiri, P.J. Keely, et al., Microtubules regulate GEF-H1 in response to extracellular matrix stiffness, *Mol. Biol. Cell* 23 (2012) 2583–2592.
- [12] E.C. Lessey, C. Guilly, K. Burrige, et al., From mechanical force to RhoA activation, *Biochemistry* 51 (2012) 7420–7432.
- [13] E. Cukierman, D.E. Bassi, et al., Physico-mechanical aspects of extracellular matrix influences on tumorigenic behaviors, *Semin. Cancer Biol.* 20 (2010) 139–145.
- [14] R.J. Petrie, N. Gavara, R.S. Chadwick, K.M. Yamada, et al., Nonpolarized signaling reveals two distinct modes of 3D cell migration, *J. Cell Biol.* 197 (2012) 439–455.
- [15] J.G. Goetz, S. Minguet, I. Navarro-Lerida, J.J. Lazcano, R. Samaniego, E. Calvo, M. Tello, T. Osteso-Ibanez, T. Pellinen, A. Echarri, A. Cerezo, A.J. Klein-Szanto, R. Garcia, P.J. Keely, P. Sanchez-Mateos, E. Cukierman, M.A. Del Pozo, et al., Biomechanical remodeling of the microenvironment by stromal caveolin-1 favors tumor invasion and metastasis, *Cell* 146 (2011) 148–163.
- [16] T. Lammermann, B.L. Bader, S.J. Monkley, T. Worbs, R. Wedlich-Soldner, K. Hirsch, M. Keller, R. Forster, D.R. Critchley, R. Fassler, M. Sixt, et al., Rapid leukocyte migration by integrin-independent flowing and squeezing, *Nature* 453 (2008) 51–55.
- [17] M. Barczyk, S. Carracedo, D. Gullberg, et al., Integrins, *Cell Tissue Res.* 339 (2010) 269–280.
- [18] P. Roca-Cusachs, N.C. Gauthier, A. Del Rio, M.P. Sheetz, et al., Clustering of alpha(5)beta(1) integrins determines adhesion strength whereas alpha(v)beta(3) and talin enable mechanotransduction, *Proc. Natl. Acad. Sci. U. S. Am.* 106 (2009) 16245–16250.
- [19] Y. Mao, J.E. Schwarzbauer, et al., Accessibility to the fibronectin synergy site in a 3D matrix regulates engagement of alpha5beta1 versus alphavbeta3 integrin receptors, *Cell Commun. Adhes.* 13 (2006) 267–277.
- [20] J.A. Askari, P.A. Buckley, A.P. Mould, M.J. Humphries, et al., Linking integrin conformation to function, *J. Cell Sci.* 122 (2009) 165–170.
- [21] E. Cukierman, R. Pankov, K.M. Yamada, et al., Cell interactions with three-dimensional matrices, *Curr. Opin. Cell Biol.* 14 (2002) 633–639.
- [22] B. Geiger, K.M. Yamada, et al., Molecular architecture and function of matrix adhesions, *Cold Spring Harbor Perspect. Biol.* 3 (2011).
- [23] A.D. Doyle, M.L. Kutys, M.A. Conti, K. Matsumoto, R.S. Adelstein, K.M. Yamada, et al., Micro-environmental control of cell migration—myosin IIA is required for efficient migration in fibrillar environments through control of cell adhesion dynamics, *J. Cell Sci.* 125 (2012) 2244–2256.
- [24] C.K. Choi, M. Vicente-Manzanares, J. Zareno, L.A. Whitmore, A. Mogilner, A.R. Horwitz, et al., Actin and alpha-actinin orchestrate the assembly and maturation of nascent adhesions in a myosin II motor-independent manner, *Nat. Cell Biol.* 10 (2008) 1039–1050.
- [25] R.J. Pelham Jr., Y. Wang, et al., Cell locomotion and focal adhesions are regulated by substrate flexibility, *Proc. Natl. Acad. Sci. U. S. Am.* 94 (1997) 13661–13665.
- [26] M.J. Paszek, N. Zahir, K.R. Johnson, J.N. Lakin, G.I. Rozenberg, A. Gefen, C.A. Reinhart-King, S.S. Margulies, M. Dembo, D. Boettiger, D.A. Hammer, V.M. Weaver, et al., Tensional homeostasis and the malignant phenotype, *Cancer Cell* 8 (2005) 241–254.
- [27] N.O. Deakin, C. Ballestrem, C.E. Turner, et al., Paxillin and Hic-5 interaction with vinculin is differentially regulated by Rac1 and RhoA, *PLoS One* 7 (2012) e37990.
- [28] J.C. Kuo, X. Han, C.T. Hsiao, J.R. Yates 3rd, C.M. Waterman, et al., Analysis of the myosin-II-responsive focal adhesion proteome reveals a role for beta-Pix in negative regulation of focal adhesion maturation, *Nat. Cell Biol.* 13 (2011) 383–393.
- [29] S.L. Gupton, C.M. Waterman-Storer, et al., Spatiotemporal feedback between actomyosin and focal-adhesion systems optimizes rapid cell migration, *Cell* 125 (2006) 1361–1374.
- [30] O. Peleg, T. Savin, G.V. Kolmakov, I.G. Salib, A.C. Balazs, M. Kroger, V. Vogel, et al., Fibers with integrated mechanochemical switches: minimalistic design principles derived from fibronectin, *Biophys. J.* 103 (2012) 1909–1918.
- [31] B.Z. Katz, E. Zamir, A. Bershadsky, Z. Kam, K.M. Yamada, B. Geiger, et al., Physical state of the extracellular matrix regulates the structure and molecular composition of cell-matrix adhesions, *Mol. Biol. Cell* 11 (2000) 1047–1060.
- [32] A.P. Mould, J.A. Askari, S. Aota, K.M. Yamada, A. Irie, Y. Takada, H.J. Mardon, M.J. Humphries, et al., Defining the topology of integrin alpha5beta1-fibronectin interactions using inhibitory anti-alpha5 and anti-beta1 monoclonal antibodies. Evidence that the synergy sequence of fibronectin is recognized by the amino-terminal repeats of the alpha5 subunit, *J. Biol. Chem.* 272 (1997) 17283–17292.
- [33] S. Aota, M. Nomizu, K.M. Yamada, et al., The short amino acid sequence Pro-His-Ser-Arg-Asn in human fibronectin enhances cell-adhesive function, *J. Biol. Chem.* 269 (1994) 24756–24761.
- [34] J.C. Friedland, M.H. Lee, D. Boettiger, et al., Mechanically activated integrin switch controls alpha5beta1 function, *Science* 323 (2009) 642–644.
- [35] P. Friedl, K. Wolf, et al., Plasticity of cell migration: a multiscale tuning model, *J. Cell Biol.* 188 (2010) 11–19.
- [36] R. Pankov, Y. Endo, S. Even-Ram, M. Araki, K. Clark, E. Cukierman, K. Matsumoto, K.M. Yamada, et al., A Rac switch regulates random versus directionally persistent cell migration, *J. Cell Biol.* 170 (2005) 793–802.
- [37] A.D. Doyle, F.W. Wang, K. Matsumoto, K.M. Yamada, et al., One-dimensional topography underlies three-dimensional fibrillar cell migration, *J. Cell Biol.* 184 (2009) 481–490.
- [38] R. Castello-Cros, D.R. Khan, J. Simons, M. Valianou, E. Cukierman, et al., Staged stromal extracellular 3D matrices differentially regulate breast cancer cell responses through PI3K and beta1-integrins, *BMC Cancer* 9 (2009) 94.
- [39] F. Berthod, L. Germain, N. Tremblay, F.A. Auger, et al., Extracellular matrix deposition by fibroblasts is necessary to promote capillary-like tube formation *in vitro*, *J. Cell Physiol.* 207 (2006) 491–498.